

Article

Relative Individual Sprint in Most Demanding Passages of Play in Spanish Professional Soccer Matches

Juan Ángel Piñero ¹, Marcos Chena ^{2,3}, Juan Carlos Zapardiel ³, Alberto Roso-Moliner ⁴,
Elena Mainer-Pardos ⁴, Miguel Lampre ^{4,*} and Demetrio Lozano ⁴

¹ CF Fuenlabrada, 28942 Madrid, Spain

² Facultad de Ciencias del Deporte de Toledo, Universidad de Castilla La Mancha, 13001 Ciudad Real, Spain

³ Faculty of Medicine and Health Sciences, Campus Universitario-C/19, University of Alcalá, Av. de Madrid, Km 33,600, 28871 Madrid, Spain

⁴ Faculty of Health Sciences, San Jorge University, Autovía A23 Km 299, 20830 Villanueva de Gállego, 50830 Zaragoza, Spain

* Correspondence: miki_tauste@hotmail.com

Abstract: (1) Background: The objective of this research was to analyse the most demanding passages (MDP) considering the sprint variable relative to the maximum level of sprint ability of each player as a function of player position, final outcome and part of the match during the competitive phase of a professional soccer season. (2) Methods: Global positioning system (GPS) data were collected from 22 players according to their playing position in the last 19 match days of the Spanish La Liga professional soccer in the 2020/2021 season. MDP were calculated from 80% of the maximum sprint speed of each player. (3) Results: Wide midfielders covered the greatest distance at >80% of the maximum speed (2.4 ± 1.63 seg) and the longest duration (21.91 ± 13.35 m) in their MDP. When the whole team was losing, it demonstrated greater distances (20.23 ± 13.04 m) and longer durations (2.24 ± 1.58 seg) compared to games in which it was winning. When the team ended up drawing, the relative sprint distance covered in the second half was significantly greater than in the first (16.12 ± 21.02 ; SD = 0.26 ± 0.28 ($-0.03/-0.54$)). (4) Conclusions: Different demands of MDP, according to the sprint variable relative to the maximum individual capacity in competition, are required when contextual game factors are considered.

Keywords: football; GPS; worst-case scenario; match analyse; sprint distance



Citation: Piñero, J.Á.; Chena, M.; Zapardiel, J.C.; Roso-Moliner, A.; Mainer-Pardos, E.; Lampre, M.; Lozano, D. Relative Individual Sprint in Most Demanding Passages of Play in Spanish Professional Soccer Matches. *Sports* **2023**, *11*, 72. <https://doi.org/10.3390/sports11040072>

Academic Editors:
Ioannis Michailidis
and Beat Knechtle

Received: 6 February 2023
Revised: 10 March 2023
Accepted: 20 March 2023
Published: 23 March 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

In modern soccer, technological advances and match and training analyses have provided coaches with scientifically based data on player performance with respect to the physical, technical and tactical dimensions of the game [1,2]. Quantifying and knowing the physical demands during competition is of great importance to better understanding what happens during competition and allowing for the adequate management of training intensity [3,4] through the adequate periodization of specific tasks. An optimal dose of training helps to conditionally prepare the soccer player for competition while minimizing the incidence of overuse injuries [5,6]. In recent years, high-intensity actions and the distance and number of sprints have increased by approximately 30–35% in official competition matches [7]. Therefore, players need to be robust enough to cope with such demands [8] and execute their tactical role effectively [9].

Moreover, to augment the intensity of the game, the rate and burden of hamstring injuries has also increased significantly since 2001, showing an average annual increase of 2.3% [10]. Although its origin is multifactorial [11], the increase in hamstring injuries could be logical as an increase in competitive demands since 70% of these injuries usually occur during sprints or high-speed runs [12]. This evolution of the game towards high-intensity, repetitive actions increases the risk of hamstring injuries [10] due to the presence

of high-force, repetitive eccentric actions [13]. Additionally, reduced recovery between these high-intensity efforts [14] may be associated with an increased risk of injury.

For this reason, control and knowledge of the most demanding passages (the most intense periods of training, or matches that show peak activities higher than the average of the match) associated with sprinting can be very useful for attempting to minimize the risk of injury and, in this way, trying to prevent hamstring injuries in soccer due to the negative impact that injuries have on soccer teams at an economic level and with respect to team performance [15–17]. Our research deals precisely with this issue, using knowledge of the most demanding passages related to individualized maximum sprint capacity to attempt to minimize or reduce the hamstring injury rate in soccer teams, allowing coaches and trainers to prescribe optimal training loads in order to prepare athletes to meet the needs of the competition.

Average physical demands, traditionally used as the main method of load quantification, could be underestimating the most demanding passages (MDP) to which players are subjected during the match [18–21] due to the intermittency of the game itself [22,23]. The MDP is the most intense period of training or matches and demonstrates peak activities that are higher than the average of the match [18,19]. The rolling average method has been one of the main instruments used to analyse the MDP [4,8,18,24,25] in different interval windows (1, 3, 5 and 10 min). The intensity of the MDP is greater when the duration of the studied period is shorter [19].

The unpredictable nature of soccer has demonstrated that the physical demands are influenced by tactical–technical factors [26]. MDP are contextually dependent, as research has shown that they can vary based on the player-specific position on the pitch [4,8,27–29], tactical role [27], play formation [30], match outcome [29], whether the team is playing at home or away [29], the first or second half of the game [4,29] and match congestion period [24].

However, despite the importance of top-speed actions in soccer, there is currently no consensus on the definition of sprinting. This is because there are different criteria regarding the defined speed thresholds in soccer [31–33]. The players' MDP is over- or underestimated depending on their sprint ability. Concerning MDP calculated with relative thresholds, the MDP values are overestimated for faster players, and underestimated for slower players [31]. To date, no known manuscript reports information on the MDP of the competition based on the maximum capacity of soccer players, which one might think is information of relative interest from a practical perspective [34]. Regarding these issues, this study has the novelty of analysing the efforts of maximum demands relative to the maximum individual sprinting capacity of each soccer player. The objective of this study was to analyse the most demanding passages (MDP), considering the sprint variable relative to the maximum level of sprint ability of each player as a function of player position, final outcome and part of the match during the competitive phase of a professional soccer season.

2. Materials and Methods

The design responds to a retrospective observational research study carried out with professional soccer players during a season. The MDP of each player was analysed depending on his maximum speed in professional soccer matches. The independent variables used were the playing position, the halftime of the match, the result of the match and the competition phase. Global positioning system (GPS) data were collected from 22 players in a total of 19 games (90 min plus extra time) from the last 19 match days in the 2020–2021 competitive season. The players provided informed consent via signature, and the study was fully approved by the Sports Management Department of the Football Club. In addition, the study complied with the research ethics standards of the University of Alcalá, Madrid, Spain, code CEI/HU/2019/08, and was conducted by the principles set out in the Declaration of Helsinki.

2.1. Participants

Twenty-two professional soccer players participated in this study (age: 25.8 ± 5 years; mass: 76.4 ± 6.5 kg; stature: 1.81 ± 0.07 m). The players belonged to a club in the Spanish La Liga professional soccer at Tier 4: the Elite/International Level [35]. Data were collected throughout 19 competitive matches in the 2020–2021 competitive season (6 wins; 6 draws; 7 losses; final position: 11th). The players were grouped according to their playing positions as central defenders (CD: $n = 4$), fullbacks (FB: $n = 4$), midfielders (MF: $n = 6$), wide midfielders (WM: $n = 4$) and forwards (FW: $n = 4$). Goalkeepers were not included in the analysis.

2.2. Research Instruments

The time motion of each player was recorded individually in all matches with a 10 Hz GPS device, as used in previous studies [36,37]. Research has shown this system to be a valid and reliable assessment for monitoring the movement demands of team players [38]. To avoid variability, each player always used the same GPS device, a WIMU PRO (RealTrackSystem, Almería, Spain), located between the two scapulae with a special vest. The same methodology used in earlier studies in which the MDP of a professional soccer game was adopted for data collection [4,8,9,24]. The coding and analysis of the data were performed with sPRO software (RealTrackSystems, Almería, Spain) [39]. To properly monitor and prescribe high-intensity training loads, predefined (absolute) or individualized (relative) criteria were used [32,33]. Although most of the published information determines the performance profile in soccer through absolute thresholds, it has been shown that these thresholds could overestimate or underestimate the amount of high- and maximum-intensity efforts of the players [32,33]. Thus, MDP were calculated from 80% of the maximum sprint speed of each player. Moreover, monitoring the distance covered for >80% of the maximum speed guarantees greater precision in the determination of sprint workloads and the associated injury risk [32]. Analysis was performed using time windows of 1 min because speeds above 80% of the maximum speed of each player are hardly bearable for longer periods [40]. This study analysed MDP during short-duration game actions. For this purpose, the range of scenarios recorded for each match was set at 50% of the maximum obtained for the selected variable, thus filtering out the cases of maximum demand scenarios in each match.

2.3. Procedures

All players performed two control tests at three months apart (September–January): the 40 m linear maximal speed test [41–43]. Maximum speed data were also recorded at every training session and match during the study (January–May) to monitor the variation in each player's maximum speed throughout the season. Each match was 90 min in duration (two 45 min halves), plus extra time (4–9 min). Data were collected based on what happened in each of the halves, the positions of the players and the result. These data were then averaged across all observations by position for the analysis between groups concerning the distance travelled and the duration of each MDP.

2.4. Data Analysis

All data are presented as means and standard deviations (mean \pm SD). Magnitude-based inferences and a precision of estimation were used to analyse the data [44]. All processed variables were log-transformed to reduce the non-uniformity of error. Differences between the halves (first half and second half) were assessed via standardised mean differences (Cohen's d) and respective 90% confidence limits. The effect size was calculated using the Cohen's d from the differences of the groups' means and the weighted standard deviation. Threshold values for standardized differences were >0.2 (small), >0.6 (moderate), >1.2 (large) and very large (>2.0) [45]. The statistical analysis was performed with the software package SPSS, version 28.0 (SPSS Inc., Chicago, IL, USA).

3. Results

Table 1 shows the mean \pm SD and confidence intervals (95%) duration (s), distance ($\text{m}\cdot\text{min}^{-1}$) and maximum speed ($\text{m}\cdot\text{s}^{-1}$) of all the MDP values for each position of the players.

Table 1. Duration, distance, and maximal running velocity to the most demanding passages according to play position.

Part of the Match	MDP									
	n	Duration (s)			Distance (m)			Vmax (m/s)		
		Values \pm SD	IC 95%	IC 95%	Values \pm SD	IC 95%	IC 95%	Values \pm SD	IC 95%	IC 95%
1° Part	151	2.05 \pm 1.37	1.83	2.27	18.98 \pm 11.42	17.15	20.82	32.15 \pm 1.63	31.89	32.41
2° Part	192	2.28 \pm 1.64	2.05	2.52	20.57 \pm 13.67	18.62	22.51	32.17 \pm 1.61	31.94	32.40

MDP: Most Demanding Passages; Vmax: Maxima velocity in competition.

Table 2 shows the mean \pm SD and confidence intervals (95%), duration (s), distance ($\text{m}\cdot\text{min}^{-1}$) and maximum running velocity ($\text{m}\cdot\text{s}^{-1}$) of all the MDP values for each half of the match.

Table 2. Duration, distance, and maximal running velocity of the most demanding passages according to part of the match.

Position	MDP									
	n	Duration (s)			Distance (m)			Vmax (m/s)		
		Values \pm SD	IC 95%	IC 95%	Values \pm SD	IC 95%	IC 95%	Values \pm SD	IC 95%	IC 95%
Central defender	75	2.15 \pm 1.48	1.81	2.49	19.26 \pm 12.01	16.47	22.04	31.74 \pm 1.79	31.33	32.15
Fullback	69	2.08 \pm 1.37	1.75	2.41	19.31 \pm 11.87	16.46	22.16	32.30 \pm 1.11	32.04	32.57
Midfielder	48	2.13 \pm 1.53	1.68	2.57	18.75 \pm 12.29	15.22	22.36	30.55 \pm 1.41	30.14	30.96
Wide midfielder	82	2.4 \pm 1.63	2.04	2.76	21.91 \pm 13.35	18.98	24.85	32.68 \pm 1.74	32.29	33.06
Forward	69	2.10 \pm 1.61	1.71	2.49	19.42 \pm 13.87	16.09	22.75	32.99 \pm 0.68	32.82	33.15

MDP: Most Demanding Passages; Vmax: Maxima velocity in competition.

Table 3 shows the mean \pm SD and confidence intervals (95%), duration (s), distance ($\text{m}\cdot\text{min}^{-1}$) and maximum speed ($\text{m}\cdot\text{s}^{-1}$) of all the MDP values according to the result of the match.

Table 3. Duration, distance, and maximal running velocity for the most demanding passages according to result of the match.

Result	MDP									
	n	Duration (s)			Distance (m)			Vmax (m/s)		
		Values \pm SD	IC 95%	IC 95%	Values \pm SD	IC 95%	IC 95%	Values \pm SD	IC 95%	IC 95%
Win	115	2.05 \pm 1.40	1.79	2.31	18.77 \pm 11.76	16.60	20.94	32.11 \pm 1.62	31.81	32.41
Draw	128	2.25 \pm 1.59	1.98	2.53	20.57 \pm 13.36	18.24	22.91	32.30 \pm 1.62	32.02	32.59
Loss	100	2.24 \pm 1.58	1.92	2.55	20.23 \pm 13.04	17.65	22.82	32.03 \pm 1.61	31.71	32.35

MDP: Most Demanding Passages; Vmax: Maxima velocity in competition.

Table 4 shows the mean \pm SD and confidence intervals (95%), duration (s), distance ($\text{m}\cdot\text{min}^{-1}$) and maximum speed ($\text{m}\cdot\text{s}^{-1}$) of all the MDP values according to the date of the season.

Table 4. Duration, distance, and maximal running velocity to the most demanding passages according to date of the season.

Date	MDP									
	n	Duration (s)			Distance (m)			Vmax (m/s)		
		Values \pm SD	IC 95%	IC 95%	Values \pm SD	IC 95%	IC 95%	Values \pm SD	IC 95%	IC 95%
J24	18	1.97 \pm 1.61	1.17	2.77	18.21 \pm 13.58	11.45	24.96	32.11 \pm 1.70	31.27	32.95

Table 4. Cont.

Date	MDP									
	n	Duration (s)			Distance (m)			Vmax (m/s)		
		Values ± SD	IC 95%		Values ± SD	IC 95%		Values ± SD	IC 95%	
J25	21	2.48 ± 1.28	1.87	3.08	22.56 ± 11.39	17.23	27.90	32.51 ± 1.73	31.70	33.32
J26	17	1.39 ± 1.01	0.85	1.93	12.73 ± 8.96	7.96	17.51	32.42 ± 1.84	31.44	33.40
J27	18	2.66 ± 1.47	1.91	3.42	23.99 ± 11.91	23.99	17.87	32.37 ± 1.75	31.37	33.27
J28	21	2.30 ± 1.32	1.70	2.90	21.62 ± 10.70	16.75	26.50	32.29 ± 1.86	31.44	33.13
J29	19	1.97 ± 1.18	1.97	2.55	17.99 ± 8.98	13.52	22.45	31.93 ± 1.56	31.15	32.71
J30	15	1.46 ± 1.12	0.81	2.11	14.09 ± 8.95	8.93	19.26	31.85 ± 1.29	31.11	31.58
J31	22	2.14 ± 1.59	2.14	1.41	20.14 ± 12.99	14.22	26.06	31.92 ± 1.49	31.24	32.60
J32	13	2.02 ± 1.60	1.00	3.03	18.81 ± 12.26	11.02	26.60	31.80 ± 1.79	30.67	32.94
J33	17	1.81 ± 1.31	1.11	2.51	16.41 ± 10.40	10.86	21.95	31.91 ± 1.34	31.20	32.63
J34	21	2.34 ± 2.10	1.39	3.30	20.58 ± 15.99	13.30	27.86	32.31 ± 1.37	31.69	32.94
J35	18	2.50 ± 1.68	1.64	3.36	22.93 ± 14.53	15.45	30.40	32.49 ± 1.34	31.80	33.18
J36	21	2.06 ± 1.38	1.41	2.71	16.74 ± 10.69	11.73	21.74	32.26 ± 1.69	31.47	33.05
J37	21	2.09 ± 1.58	1.33	2.86	19.95 ± 13.57	13.41	26.49	32.24 ± 1.59	31.47	33.00
J38	19	1.79 ± 1.25	1.19	2.40	16.19 ± 9.54	11.59	20.79	32.17 ± 1.54	31.43	32.91
J39	17	3.58 ± 1.92	2.56	4.61	31.95 ± 16.87	22.96	40.94	31.91 ± 1.76	30.97	32.85
J40	20	2.30 ± 1.66	1.50	3.10	20.66 ± 13.76	14.03	27.29	32.34 ± 1.84	31.46	33.23
J41	19	2.54 ± 1.56	1.76	3.32	23.22 ± 14.83	15.84	30.59	32.18 ± 1.67	31.35	33.01
J42	22	1.90 ± 1.33	1.30	2.51	17.78 ± 11.47	12.55	23.00	31.85 ± 1.88	31.00	32.71

MDP: Most Demanding Passages; Vmax: Maxima velocity in competition.

Table 5 shows the difference standardized (Cohen) of the MDP according to position on the half of the match.

Table 5. Difference standardized (Cohen) to the Most Demanding Passages according half of the mach.

Position	MDP	1st Half	2nd Half	Difference (%)	Difference Standardized (Cohen)	Chances	Qualitative
Central defender	Distance (m)	15.36 ± 10.54 (36)	22.30 ± 12.77 (39)	31.14 ± 17.46	0.59 ± 0.38 (0.21/0.96)	95/5/0	Very Likely
	Duration (s)	1.77 ± 1.26 (36)	2.48 ± 1.59 (39)	28.43 ± 20.52	0.49 ± 0.38 (0.11/0.86)	89/10/0	Likely
Fullback	Distance (m)	21.17 ± 11.03 (39)	16.40 ± 1.41 (33)	29.12 ± 13.24	-0.40 ± 0.39 (-0.78/-0.01)	1/19/80	Likely
	Duration (s)	2.28 ± 1.33 (39)	2.19 ± 1.83 (33)	20.62 ± 5.28	-0.28 ± 0.40 (-0.68/0.12)	2/34/63	Possibly
Midfielder	Distance (m)	15.53 ± 10.47 (26)	17.44 ± 14.90 (22)	10.93 ± 29.72	0.15 ± 0.44 (-0.30/0.59)	42/48/10	Unclear
	Duration (s)	1.94 ± 1.16 (26)	2.29 ± 1.81 (22)	15.21 ± 35.95	0.23 ± 0.47 (-0.25/0.70)	54/40/7	Unclear
Wide midfielder	Distance (m)	20.00 ± 14.23 (32)	21.18 ± 13.67 (51)	5.56 ± 4.14	0.08 ± 0.37 (-0.29/0.45)	30/60/10	Unclear
	Duration (s)	2.28 ± 1.70 (32)	2.47 ± 1.60 (51)	7.94 ± 6.42	0.12 ± 0.38 (-0.26/0.50)	36/56/8	Unclear
Forward	Distance (m)	19.25 ± 11.33 (30)	19.55 ± 15.69 (39)	1.51 ± 27.77	0.02 ± 0.39 (-0.37/0.41)	22/60/18	Unclear
	Duration (s)	1.98 ± 1.31 (30)	2.19 ± 1.83 (39)	9.94 ± 28.68	0.14 ± 0.39 (-0.26/0.53)	39/53/8	Unclear

MDP: Most Demanding Passages.

Table 6 shows the difference standardized (Cohen) of the MDP according to the result of the match on the half of the match.

Table 6. Difference standardized (Cohen) of the most demanding passages according to result of the match and the half of the match.

Results	MDP	1st Half	2nd Half	Difference (%)	Difference Standardized (Cohen)	Chances	Qualitative
Win	Distance (m)	17.56 ± 11.43 (54)	17.93 ± 12.76 (66)	2.11 ± 10.44	0.03 ± 0.30 (-0.27/0.33)	17/72/10	Unclear
	Duration (s)	2.03 ± 1.29 (54)	2.07 ± 1.50 (66)	2.24 ± 14.03	0.03 ± 0.31 (0.27/0.34)	18/71/11	Unclear
Draw	Distance (m)	18.02 ± 11.65 (56)	21.48 ± 14.76 (74)	16.12 ± 21.02	0.26 ± 0.28 (-0.03/-0.54)	63/36/0	Possibly
	Duration (s)	2.01 ± 1.38 (39)	14.76 ± 1.72 (74)	17.89 ± 19.61	0.28 ± 0.29 (-0.01/0.57)	67/32/0	Possibly
Loss	Distance (m)	19.56 ± 12.31 (45)	19.19 ± 14.14 (59)	1.93 ± 12.99	-0.03 ± 0.31 (-0.35/0.29)	12/69/19	Unclear
	Duration (s)	2.14 ± 1.46 (45)	2.32 ± 1.69 (59)	8.12 ± 13.52	0.12 ± 0.33 (-0.21/0.45)	34/60/6	Unclear

MDP: Most Demanding Passages.

Figure 1 shows effect size on the MDP according to the position and half of the match.




Figure 1. Effect size to the Most Demanding Passages according to position and half of the match.

Figure 2 shows the effect size on the MDP according to the result of the match.




Figure 2. Effect size on the most demanding passages according to the result of the match.

4. Discussion

The objectives of the present study were: (i) to analyse the MDP, considering the sprint variable relative to the maximum level of sprint ability of each player during the competitive phase of a professional soccer season; (ii) to compare the MDP of each playing position; and (iii) to compare the MDP between the first and second halves of each match.

This is the first study to analyse the efforts of maximum demands relative to the maximum individual capacities of each soccer player.

Competitive demands have led coaches and physical trainers to challenge the principles of training through the prescription of optimal stimuli to develop or maintain adaptations without exceeding the limits of physiological tolerance of the players [36,46]. For this purpose, the MDP must be considered in the training context since planning according to the average competition references may not be a sufficient stimulus to prepare soccer players for the most demanding phases [8,47]. To describe the MDP, the calculation of the moving average has been considered the most accurate method for determining the most intense periods [8,34], as it was calculated in this study. However, most authors have focused their study on describing the MDP in terms of the variables total distance per minute, distance covered at a high intensity (>19.8 km/h) per minute and distance covered at a sprint (>25.2 km/h) per minute in periods of 1, 3, 5 and 10 min [8,27,28,34].

Unlike the works mentioned above, this study attempted to analyse the sprint MDP by considering sprint the ability to make a maximum effort or very close to the maximum individual capacity of each player to obtain the distance, duration time and speed information in those competitive situations. Of the locomotor requirements demanded by the game, the sprint is probably one of the most determining since it precedes most scoring opportunities and is the most frequent mechanism of muscle injury.

Players have been shown to reach 85–95% of their maximum speed during soccer matches [46]. Therefore, it appears that training conditions of maximal stress based on an individual soccer player's ability to optimally meet competitive demands is required [46,48,49]. On the other hand, O'Connor et al. (2020) [32] demonstrated significant differences in the results obtained when they analysed the relationship between the sprint workload, analysed with relative and absolute thresholds, and the incidence of injuries, concluding that monitoring the distance covered at >80% of the maximum speed guarantees greater precision in the determination of sprint workloads and associated injury risk. Accordingly, and considering that the fitness of athletes is variable throughout the competitive period, in this study, it was decided to control the sprint variable using the relative threshold of >80% of the maximum individual speed reached in competition, which was automatically updated to the maximum value obtained throughout the season. However, in some cases, such as for players with a certain time of inactivity or low participation and poor competitive rhythm, this value of maximum speed obtained at a specific moment would be higher than the real fitness that these players have for the speed. For this reason, a relative range of 80% of the maximum sprint was placed for those players whose top speed was overestimated due to their low fitness to determinate a more accurate MDP value relative to the maximum speed at this particular moment, guaranteeing greater precision in the determination of sprint workloads and the associated injury risk.

This study showed that during the MDP in the matches, there were differences between the different positions, as observed in previous studies [8,18,29]. The results showed that wide midfielders were the players who covered the greatest distance at >80% of the maximum speed in their MDP and demonstrated the longest duration of these efforts. According to the scientific literature, wide midfielders undergo greater distances at a high intensity (HSR) during competition [26,50–54]. Nonetheless, Martín-García and Gómez Díaz et al. (2018) [8] did not find the highest HSR values in the MDP of these players. These authors warned that their results could have been different if they had used the HSR as a criterion variable, as it was used in other publications [18]. However, the maximum capacities of the athletes were not taken into account in any of the studies that employed HSR as a criterion [33], registering said variable with absolute criteria.

Although midfielders must cover great distances throughout a game [26,51,52,54], when the total distance was analysed regardless of speed, these players were the ones who covered the most meters per minute [8]. In this study, it was observed that according to their MDP, the distance they covered in the relative sprint was less than other positions. Therefore, these results confirm that midfielders are players with a high level of intervention

in game actions and are capable of recovering great distances but do not have the need to reach the maximum demands of their capacity [51].

In the analysis of the defensive line, there were differences in the MDP regarding the distance covered in a relative sprint between halves. Fullbacks covered more in the first part than the second; however, central defenders covered a significantly greater distance and duration relative to the sprint in the second half. In line with these results, Rey et al. (2020) reported that central defenders increased the sprint distance covered during the second half of the match [55]. Soccer matches tend to present less control and higher transition situations at the end of the matches due to the need to win. This could explain the fact that central defenders covered greater distances and durations relative to their sprint, since central defenders demonstrate most of their high intensity actions in recovery runs and coverage [56], defensive situations that occur constantly in transition situations. In our study, no statistically significant differences were found between both parties in the rest of the positions (wide midfielder and forward).

Moreover, analysing the competitive physical demands of soccer players based on the total playing time could overestimate fatigue-induced decreases in performance. Rey et al. (2020) [55] found that there were no significant differences when the whole team is taken as a reference, without differentiating between positions, in the efforts made by the professional soccer players between the first and the second parts, taking into account the useful time of exposure. However, it seems that the physical demands could be positionally dependent, since when we differentiate between positions, central defenders and central midfielders increased the sprint distance covered during the second half of the match [55]. Therefore, the effective playing time and playing position should be taken into account when analysing the physical demands and performance of soccer players.

Regarding the relationship with the final score, Oliva-Lozano et al. (2020) [29] exposed that the result was a contextual variable with a significant impact on the most demanding competitive scenarios in total distance, high-speed running and sprint distance in the periods of 1 and 3 min. Winning the match resulted in higher values of total distance covered, the distance at high intensity and the distance at sprint compared to drawing and losing the game [29]. Diez et al. (2021) [57] also demonstrated that more SPR are exhibited by players when the team plays at home and wins. However, in contrast to these studies, our results indicated that games in which the team was losing showed greater distances and longer durations in maximum efforts compared to games in which it was winning. The timing of the matches selected in this study included the last match weeks of the season (J19 to J42), in which the teams fight for their objectives and each point won or lost has greater relevance. Perhaps this would justify the results of our study, since when teams are losing, they need to make greater physical efforts to reduce that difference and try to win. Moreover, in line with these results, when considering the MDP of the matches in which the team ended up drawing, the relative sprint distance covered in the second half was significantly greater than in the first, perhaps due to the need to score to achieve the classifying objectives. Nobari et al. (2021) [58] also highlighted the difference between drawn versus won matches, explaining that the similar performance may be related to the duration of the match and therefore the intensity generated in both.

For the whole team, a greater duration and distance for the MDP of the sprint covered in the second part could have an impact on hamstring injuries since the hamstring fatigue resistance has been shown to be a soccer-specific risk factor for this type of injury [59,60]. In line, Raya et al. (2020) [61] reported that the 76–90 min time period presented the highest values of hamstring injury: 12 out of 63 injuries (19.04% of total) in comparison to all selected time periods in an Elite Spanish Male Academy. Sprinting is a key area for improving performance and injury prevention. The authors of [46,50,62] showed that professional soccer players who were trained to a higher chronic training load were able to better tolerate a weekly acute increase in exposure to maximum speed events as a consequence of the protective effect of chronic, high training loads. Therefore, these data suggest the need for athletes to regularly expose themselves during training to maximum

speeds [32,46] and in fatigue conditions to prepare for competition. This could help coaches plan and structure their training and competitions and try to minimize or reduce the injury rate.

Future research on the relationship between sprinting at relative thresholds and MDP linked to training tasks that can replicate competitive demands seems necessary, with the aim of optimizing the training process to improve performance and prevent injuries.

5. Conclusions

The physical performance profile in soccer should not be understood solely from the average analysis of competitive demands, since this could be hiding the maximum demands covered by players. Players do not show a decrease in physical performance as the match progresses, even increasing their performance in situations of maximum demand in the second half. Moreover, other contextual factors as the result of the match, playing position and the moment of the season can influence the MDP.

Coaches and trainers must prescribe optimal sprint training loads, prioritizing both quality and quantity in situations of fatigue, due to the increased duration and distance in the MDP of the sprint covered in the second half, as demonstrated by this study, in order to prepare athletes for meeting competitive needs and to reduce the risk of hamstring injury. Moreover, these tasks should be oriented to individual and positional needs, due to the tactical and positional dependence on physical demands.

It is therefore necessary to detect which are the periods of maximum demand during the matches and which of them can be generated through specific training tasks to replicate the demands of the competition by taking the athlete to situations at the limit of his capacity.

The main practical application of knowing these demands is the ability to apply and practice training tasks that meet these needs, both individually and positionally, on a day-to-day basis, not only to improve performance but also to have a protective effect on the relationship that sprinting has been demonstrated to have with both.

Author Contributions: Conceptualization, J.Á.P. and D.L.; methodology, J.Á.P., M.C., A.R.-M. and J.C.Z.; validation, M.L., E.M.-P. and D.L.; formal analysis, J.Á.P., M.C. and D.L.; investigation, J.Á.P., M.C. and D.L.; data curation, J.Á.P., M.C. and D.L.; writing—original draft preparation, J.Á.P., M.C., A.R.-M. and J.C.Z.; writing—review and editing, M.L. and J.C.Z.; supervision, A.R.-M. and D.L.; project administration, D.L. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: The study was conducted according to the guidelines of the Declaration of Helsinki and approved by the Ethics Committee of Universidad de Alcalá (Madrid, Spain) No. 31/2021.

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: Not applicable.

Acknowledgments: Research Group ValorA, under grant No. S08_20R, and Fondo Social Europeo “Construyendo Europa desde Aragón”.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Sarmiento, H.; Marcelino, R.; Anguera, M.T.; Campaniço, J.; Matos, N.; Leitão, J.C.; Leitao, J.C. Match analysis in football: A systematic review. *J. Sports Sci.* **2014**, *32*, 1831–1843. [\[CrossRef\]](#)
2. Collet, C. The possession game? A comparative analysis of ball retention and team success in European and international football, 2007–2010. *J. Sports Sci.* **2013**, *31*, 123–136. [\[CrossRef\]](#)
3. Cohen, S.B.; Towers, J.D.; Zoga, A.; Irrgang, J.J.; Makda, J.; Deluca, P.F.; Bradley, J.P. Hamstring injuries in professional football players: Magnetic resonance imaging correlation with return to play. *Sports Health* **2011**, *3*, 423–430. [\[CrossRef\]](#)
4. Casamichana, D.; Castellano, J.; Diaz, A.G.; Gabbett, T.J.; Martin-Garcia, A. The most demanding passages of play in football competition: A comparison between halves. *Biol. Sport* **2019**, *36*, 233–240. [\[CrossRef\]](#)

5. Akenhead, R.; Nassis, G.P. Training load and player monitoring in high-level football: Current practice and perceptions. *Int. J. Sports Physiol. Perform.* **2016**, *11*, 587–593. [[CrossRef](#)]
6. Weston, M. Training load monitoring in elite English soccer: A comparison of practices and perceptions between coaches and practitioners. *Sci. Med. Footb.* **2018**, *2*, 216–224. [[CrossRef](#)]
7. Barnes, C.; Archer, D.T.; Hogg, B.; Bush, M.; Bradley, P.S. The evolution of physical and technical performance parameters in the english premier league. *Int. J. Sports Med.* **2014**, *35*, 1095–1100. [[CrossRef](#)] [[PubMed](#)]
8. Martín-García, A.; Gómez Díaz, A.; Bradley, P.S.; Morera, F.; Casamichana, D. Quantification of a professional football team's external load using a microcycle structure. *J. Strength Cond. Res.* **2018**, *32*, 3511–3518. [[CrossRef](#)]
9. Lacome, M.; Simpson, B.M.; Cholley, Y.; Lambert, P.; Buchheit, M. Small-sided games in elite soccer: Does one size fit all? *Int. J. Sports Physiol. Perform.* **2018**, *13*, 568–576. [[CrossRef](#)]
10. Ekstrand, J.; Waldén, M.; Häggglund, M. Hamstring injuries have increased by 4% annually in men's professional football, since 2001: A 13-year longitudinal analysis of the UEFA Elite Club injury study. *Br. J. Sports Med.* **2016**, *50*, 731–737. [[CrossRef](#)]
11. Meeuwisse, W.H.; Tyreman, H.; Hagel, B.; Emery, C. A dynamic model of etiology in sport injury: The recursive nature of risk and causation. *Clin. J. Sport Med.* **2007**, *17*, 215–219. [[CrossRef](#)] [[PubMed](#)]
12. Ekstrand, J.; Healy, J.C.; Waldén, M.; Lee, J.C.; English, B.; Häggglund, M. Hamstring muscle injuries in professional football: The correlation of MRI findings with return to play. *Br. J. Sports Med.* **2012**, *46*, 112–117. [[CrossRef](#)]
13. Opar, D.A.; Williams, M.D.; Timmins, R.G.; Hickey, J.; Duhig, S.J.; Shield, A.J. Eccentric hamstring strength and hamstring injury risk in Australian footballers. *Med. Sci. Sports Exerc.* **2015**, *47*, 857–865. [[CrossRef](#)] [[PubMed](#)]
14. Carling, C.; Gall, F.L.; Reilly, T.P. Effects of physical efforts on injury in elite soccer. *Int. J. Sports Med.* **2010**, *31*, 180–185. [[CrossRef](#)]
15. Ekstrand, J. Preventing injuries in professional football: Thinking bigger and working together. *Br. J. Sports Med.* **2016**, *50*, 709–710. [[CrossRef](#)] [[PubMed](#)]
16. Ekstrand, J.; Askling, C.; Magnusson, H.; Mithoefer, K. Return to play after thigh muscle injury in elite football players: Implementation and validation of the Munich muscle injury classification. *Br. J. Sports Med.* **2013**, *47*, 769–774. [[CrossRef](#)]
17. Carling, C.; McCall, A.; Le Gall, F.; Dupont, G. The impact of short periods of match congestion on injury risk and patterns in an elite football club. *Br. J. Sports Med.* **2016**, *50*, 764–768. [[CrossRef](#)]
18. Delaney, J.A.; Thornton, H.R.; Rowell, A.E.; Dascombe, B.J.; Aughey, R.J.; Duthie, G.M. Modelling the decrement in running intensity within professional soccer players. *Sci. Med. Footb.* **2018**, *2*, 86–92. [[CrossRef](#)]
19. Martín-García, A.; Casamichana, D.; Gómez Díaz, A.; Cos, F.; Gabbett, T.J. Positional differences in the most demanding passages of play in football competition. *J. Sport. Sci. Med.* **2018**, *17*, 563–570.
20. Gabbett, T.J.; Kennelly, S.; Sheehan, J.; Hawkins, R.; Milsom, J.; King, E.; Whiteley, R.; Ekstrand, J. If overuse injury is a 'training load error', should undertraining be viewed the same way? *Br. J. Sports Med.* **2016**, *50*, 1017–1018. [[CrossRef](#)]
21. Mernagh, D.; Weldon, A.; Wass, J.; Phillips, J.; Parmar, N.; Waldron, M.; Turner, A. A comparison of match demands using ball-in-play versus whole match data in professional soccer players of the english championship. *Sports* **2021**, *9*, 76. [[CrossRef](#)] [[PubMed](#)]
22. Rampinini, E.; Impellizzeri, F.M.; Castagna, C.; Abt, G.; Chamari, K.; Sassi, A.; Marcora, S.M. Factors influencing physiological responses to small-sided soccer games. *J. Sports Sci.* **2007**, *25*, 659–666. [[CrossRef](#)] [[PubMed](#)]
23. Di Salvo, V.; Baron, R.; Tschan, H.; Calderon Montero, F.J.; Bachl, N.; Pigozzi, F. Performance characteristics according to playing position in elite soccer. *Int. J. Sports Med.* **2007**, *28*, 222–227. [[CrossRef](#)] [[PubMed](#)]
24. Castellano, J.; Martín-García, A.; Casamichana, D. Most running demand passages of match play in youth soccer congestion period. *Biol. Sport* **2020**, *37*, 367–373. [[CrossRef](#)] [[PubMed](#)]
25. Varley, M.C.; Fairweather, I.H.; Aughey, R.J. Validity and reliability of GPS for measuring instantaneous velocity during acceleration, deceleration, and constant motion. *J. Sports Sci.* **2012**, *30*, 121–127. [[CrossRef](#)]
26. Bradley, P.S.; Carling, C.; Gomez Diaz, A.; Hood, P.; Barnes, C.; Ade, J.; Boddy, M.; Krstrup, P.; Mohr, M. Match performance and physical capacity of players in the top three competitive standards of English professional soccer. *Hum. Mov. Sci.* **2013**, *32*, 808–821. [[CrossRef](#)]
27. Ju, W.; Doran, D.A.; Hawkins, R.; Gómez-Díaz, A.; Martín-García, A.; Ade, J.; Laws, A.; Evans, M.; Bradley, P. Contextualised peak periods of play in English Premier League matches. *Biol. Sport* **2021**, *39*, 973–983. [[CrossRef](#)]
28. Martín-García, A.; Castellano, J.; Díaz, A.G.; Cos, F.; Casamichana, D. Positional demands for various-sided games with goalkeepers according to the most demanding passages of match play in football. *Biol. Sport* **2019**, *36*, 171–180. [[CrossRef](#)]
29. Oliva-Lozano, J.M.; Rojas-Valverde, D.; Gómez-Carmona, C.D.; Fortes, V.; Pino-Ortega, J. Worst case scenario match analysis and contextual variables in professional soccer players: A longitudinal study. *Biol. Sport* **2020**, *37*, 429–436. [[CrossRef](#)]
30. Riboli, A.; Semeria, M.; Coratella, G.; Esposito, F. Effect of formation, ball in play and ball possession on peak demands in elite soccer. *Biol. Sport* **2021**, *38*, 195–205. [[CrossRef](#)]
31. Murray, N.B.; Gabbett, T.J.; Townshend, A.D. The use of relative speed zones in Australian football: Are we really measuring what we think we are? *Int. J. Sports Physiol. Perform.* **2018**, *13*, 442–451. [[CrossRef](#)] [[PubMed](#)]
32. O'Connor, F.; Thornton, H.R.; Ritchie, D.; Anderson, J.; Bull, L.; Rigby, A.; Leonard, Z.; Stern, S.; Bartlett, J.D. Greater association of relative thresholds than absolute thresholds with noncontact lower-body injury in professional Australian rules footballers: Implications for sprint monitoring. *Int. J. Sports Physiol. Perform.* **2020**, *15*, 204–212. [[CrossRef](#)] [[PubMed](#)]

33. Sweeting, A.J.; Cormack, S.J.; Morgan, S.; Aughey, R.J. When is a sprint a sprint? A review of the analysis of team-sport athlete activity profile. *Front. Physiol.* **2017**, *8*, 432. [[CrossRef](#)] [[PubMed](#)]
34. Oliva-Lozano, J.M.; Martín-Fuentes, I.; Fortes, V.; Muyor, J.M. Differences in worst-case scenarios calculated by fixed length and rolling average methods in professional soccer match-play. *Biol. Sport* **2021**, *38*, 325. [[CrossRef](#)]
35. McKay, A.K.A.; Stellingwerff, T.; Smith, E.S.; Martin, D.T.; Mujika, I.; Goosey-Tolfrey, V.L.; Sheppard, J.; Burke, L.M. Defining training and performance caliber: A participant classification framework. *Int. J. Sports Physiol. Perform.* **2022**, *17*, 317–331. [[CrossRef](#)]
36. Chena, M.; Morcillo, J.A.; Rodríguez-Hernández, M.L.; Zapardiel, J.C.; Owen, A.; Lozano, D. The Effect of Weekly Training Load across a Competitive Microcycle on Contextual Variables in Professional Soccer. *Int. J. Environ. Res. Public Health* **2021**, *18*, 5091. [[CrossRef](#)] [[PubMed](#)]
37. Di Prampero, P.E.; Botter, A.; Osgnach, C. The energy cost of sprint running and the role of metabolic power in setting top performances. *Eur. J. Appl. Physiol.* **2015**, *115*, 451–469. [[CrossRef](#)] [[PubMed](#)]
38. Hoppe, M.W.; Baumgart, C.; Polglaze, T.; Freiwald, J. Validity and reliability of GPS and LPS for measuring distances covered and sprint mechanical properties in team sports. *PLoS ONE* **2018**, *13*, e0192708. [[CrossRef](#)]
39. Pino-Ortega, J.; Rico-González, M.; Gantois, P.; Nakamura, F.Y. Level of agreement between sPRO and Kubios software in the analysis of RR intervals obtained by a chest strap. *J. Sport. Eng. Technol.* **2021**, *9*, 1–6. [[CrossRef](#)]
40. Riboli, A.; Esposito, F.; Coratella, G. The distribution of match activities relative to the maximal intensities in elite soccer players: Implications for practice. *Res. Sport. Med.* **2022**, *30*, 463–474. [[CrossRef](#)]
41. Pardos-Mainer, E.; Casajús, J.A.; Gonzalo-Skok, O. Reliability and sensitivity of jumping, linear sprinting and change of direction ability tests in adolescent female football players. *Sci. Med. Footb.* **2018**, *3*, 183–190. [[CrossRef](#)]
42. Sassi, R.H.; Dardouri, W.; Yahmed, M.H.; Gmada, N.; Mahfoudhi, M.E.; Gharbi, Z. Relative and absolute reliability of a modified agility T-test and its relationship with vertical jump and straight sprint. *J. Strength Cond. Res.* **2009**, *23*, 1644–1651. [[CrossRef](#)] [[PubMed](#)]
43. Munro, A.G.; Herrington, L.C. Between-session reliability of four hop tests and the agility T-test. *J. Strength Cond. Res.* **2011**, *25*, 1470–1477. [[CrossRef](#)] [[PubMed](#)]
44. Batterham, A.M.; Hopkins, W.G. Making meaningful inferences about magnitudes. *Int. J. Sports Physiol. Perform.* **2006**, *1*, 50–57. [[CrossRef](#)]
45. Hopkins, W.G.; Marshall, S.W.; Batterham, A.M.; Hanin, J. Progressive statistics for studies in sports medicine and exercise science. *Med. Sci. Sports Exerc.* **2009**, *41*, 3–13. [[CrossRef](#)] [[PubMed](#)]
46. Malone, S.; Owen, A.; Mendes, B.; Hughes, B.; Collins, K.; Gabbett, T.J. High-speed running and sprinting as an injury risk factor in soccer: Can well-developed physical qualities reduce the risk? *J. Sci. Med. Sport* **2018**, *21*, 257–262. [[CrossRef](#)] [[PubMed](#)]
47. Gabbett, T.J.; Hulin, B.T.; Blanch, P.; Whiteley, R. High training workloads alone do not cause sports injuries: How you get there is the real issue. *Br. J. Sports Med.* **2016**, *50*, 444–445. [[CrossRef](#)] [[PubMed](#)]
48. Mendiguchia, J.; Conceição, F.; Edouard, P.; Fonseca, M.; Pereira, R.; Lopes, H.; Morin, J.-B.; Jiménez-Reyes, P. Sprint versus isolated eccentric training: Comparative effects on hamstring architecture and performance in soccer players. *PLoS ONE* **2020**, *15*, e0228283. [[CrossRef](#)]
49. Mendiguchia, J.; Castaño-Zambudio, A.; Jiménez-Reyes, P.; Morin, J.; Edouard, P.; Conceição, F.; Tawiah-Dodoo, J.; Colyer, S.L. Can We Modify Maximal Speed Running Posture? Implications for Performance and Hamstring Injury Management. *Int. J. Sports Physiol. Perform.* **2021**, *1*, 374–383. [[CrossRef](#)]
50. Ade, J.; Fitzpatrick, J.; Bradley, P.S. High-intensity efforts in elite soccer matches and associated movement patterns, technical skills and tactical actions. Information for position-specific training drills. *J. Sports Sci.* **2016**, *34*, 2205–2214. [[CrossRef](#)]
51. Bradley, P.S.; Ade, J.D. Are current physical match performance metrics in elite soccer fit for purpose or is the adoption of an integrated approach needed? *Int. J. Sports Physiol. Perform.* **2018**, *13*, 656–664. [[CrossRef](#)] [[PubMed](#)]
52. Bradley, P.S.; Carling, C.; Archer, D.; Roberts, J.; Dodds, A.; Di Mascio, M.; Paul, D.; Gomez Diaz, A.; Peart, D.; Krstrup, P. The effect of playing formation on high-intensity running and technical profiles in English FA Premier League soccer matches. *J. Sports Sci.* **2011**, *29*, 821–830. [[CrossRef](#)]
53. Carling, C.; Le Gall, F.; Dupont, G. Analysis of repeated high-intensity running performance in professional soccer. *J. Sports Sci.* **2012**, *30*, 325–336. [[CrossRef](#)] [[PubMed](#)]
54. Castellano, J.; Alvarez-Pastor, D.; Bradley, P.S. Evaluation of Research Using Computerised Tracking Systems (Amisco[®] and Prozone[®]) to Analyse Physical Performance in Elite Soccer: A Systematic Review. *Sport. Med.* **2014**, *44*, 701–712. [[CrossRef](#)] [[PubMed](#)]
55. Rey, E.; Kalén, A.; Lorenzo-Martínez, M.; López-Del Campo, R.; Nevado-Garrosa, F.; Lago-Peñas, C. Elite Soccer Players do Not Cover Less Distance in the Second Half of the Matches When Game Interruptions Are Considered. *J. Strength Cond. Res.* **2020**, *17*. [[CrossRef](#)]
56. Ju, W.; Doran, D.; Hawkins, R.; Evans, M.; Laws, A.; Bradley, P. Contextualised high-intensity running profiles of elite football players with reference to general and specialised tactical roles. *Biol. Sport* **2023**, *40*, 291–301. [[CrossRef](#)] [[PubMed](#)]
57. Díez, A.; Lozano, D.; Arjol-Serrano, J.L.; Mainer-Pardos, E.; Castillo, D.; Torrontegui-Duarte, M.; Nobari, H.; Jaén-Carrillo, D.; Lampre, M. Influence of contextual factors on physical demands and technical-tactical actions regarding playing position in professional soccer players. *BMC Sports Sci. Med. Rehabil.* **2021**, *13*, 157. [[CrossRef](#)]

58. Nobari, H.; Oliveira, R.; Brito, J.P.; Pérez-gómez, J.; Clemente, F.M.; Ardigò, L.P. Comparison of running distance variables and body load in competitions based on their results: A full-season study of professional soccer players. *Int. J. Environ. Res. Public Health* **2021**, *18*, 2077. [[CrossRef](#)]
59. Greig, M. The influence of soccer-specific fatigue on peak isokinetic torque production of the knee flexors and extensors. *Am. J. Sports Med.* **2008**, *36*, 1403–1409. [[CrossRef](#)]
60. Buckthorpe, M.; Wright, S.; Bruce-Low, S.; Nanni, G.; Sturdy, T.; Gross, A.S.; Bowen, L.; Styles, B.; Della Villa, S.; Davison, M.; et al. Recommendations for hamstring injury prevention in elite football: Translating research into practice. *Br. J. Sports Med.* **2019**, *53*, 449–456. [[CrossRef](#)]
61. Raya-González, J.; Croix, M. de S.; Read, P.; Castillo, D. A longitudinal investigation of muscle injuries in an elite spanish male academy soccer club: A hamstring injuries approach. *Appl. Sci.* **2020**, *10*, 1610. [[CrossRef](#)]
62. Caldbeck, P.M. *Contextual Sprinting in Premier League Football*; Liverpool John Moores University (United Kingdom): Liverpool, UK, 2020; ISBN 9798582541455.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.